



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

Solution by the PROPOSER.

$r(a^2 \sin^2 \theta + b^2 \cos^2 \theta) = 2ab^2 \cos \theta$ or $u = \frac{a^2 \sin^2 \theta + b^2 \cos^2 \theta}{2ab^2 \cos \theta}$ is polar equation to the ellipse with vertex as pole.

$F = h^2 u^2 \left(\frac{d^2 u}{d\theta^2} + u \right)$ is the force required, h being an undetermined constant.

$$\frac{d^2 u}{d\theta^2} = \frac{(a^2 - b^2) \cos^4 \theta + a^2 \sin^2 \theta + a^2}{2ab^2 \cos^3 \theta}; \quad \frac{d^2 u}{d\theta^2} + u = \frac{a}{b^2 \cos^3 \theta}.$$

$$\therefore F = \frac{ah^2 u^2}{b^2 \cos^3 \theta} = \frac{ah^2 (a^2 \sin^2 \theta + b^2 \cos^2 \theta)^2}{4a^2 b^6 \cos^5 \theta},$$

$$\therefore F = \frac{a^3 h^2}{4b^6 \cos^5 \theta} - \frac{h^2 a(a^2 - b^2)}{2b^6 \cos^3 \theta} + \frac{(a^2 - b^2)^2 h^2}{4ab^6 \cos \theta}.$$

Also solved by J. SCHEFFER.

99. Proposed by F. P. MATZ, Sc. D., Ph. D., Professor of Mathematics and Astronomy in Defiance College, Defiance, Ohio.

Through the zenith of an observer at the sea-coast in north latitude $\varphi = 40^\circ$, a "cat's-tail cloud," height $h = 10$ miles, extends northeast until it touches the horizon. How far from the observer is the *advance-end* of the cloud? What is the length of the cloud measured from the end specified to the observer's zenith?

Solution by G. B. M. ZERR, A. M., Ph. D., Professor of Chemistry and Physics, The Temple College, Philadelphia, Pa.

Regarding the earth as a perfect sphere of radius 3963 miles, and disregarding the observer's height, the distance and length required will be the same for any latitude.

Let C be the center of the earth, O the position of the observer, Z his zenith, and BOA the diameter of the visible horizon.

$OA = \sqrt{(CA^2 - CO^2)} = \sqrt{[(3973)^2 - (3963)^2]} = 281.71$ miles, the distance of advance end from observer.

$$\cos OCA = \frac{3963\frac{2}{3}}{3973\frac{1}{3}} = .997483. \quad \angle OCA = 4^\circ 4'.$$

$$\therefore ZA = (4\frac{1}{15}\pi \times 7946)/360 = 281.99 \text{ miles, length of cloud.}$$

100. Proposed by F. P. MATZ, Sc. D., Ph. D., Professor of Mathematics and Astronomy in Defiance College, Defiance, O.

Determine the maximum value of $(\varphi - \varphi')$, if given electric currents C and C' produce deflections φ and φ' in a tangent galvanometer, so that $\tan \varphi / \tan \varphi' = C / C'$.

Solution by GEORGE R. DEAN, Professor of Mathematics, Missouri State School of Mines and Metallurgy, Rolla, Mo.

$$\text{We have } C' \tan \varphi = C \tan \varphi', \quad u = \varphi - \varphi'.$$

Differentiating, we must have for a maximum or minimum, $C'\sec^2\varphi d\varphi = C\sec^2\varphi'd\varphi'$. $d\varphi = d\varphi'$.

Eliminating the differentials, $C'\sec^2\varphi = C\sec^2\varphi'$.

Solving this and the given relation for φ , we find

$$\tan^2\varphi = \frac{C^2 - CC'}{C'^2 - CC'} = -\frac{C}{C'}.$$

$$\text{But } \frac{-C}{C'} = -\frac{\tan\varphi}{\tan\varphi'}. \quad \text{Hence } \tan\varphi\tan\varphi' = -1.$$

This shows that $\varphi - \varphi' = 90^\circ$.

Also solved by *H. C. WHITAKER*, and *G. B. M. ZERR*.

PROBLEMS FOR SOLUTION.

ALGEBRA.

145. Proposed by *JOHN M. COLAW*, A. M., Monterey, Va.

Solve the equations:

$$\begin{aligned} x + y + z + u + w &= 1, \\ ax + by + cz + du + ew &= h, \\ a^2x + b^2y + c^2z + d^2u + e^2w &= h^2, \\ a^3x + b^3y + c^3z + d^3u + e^3w &= h^3, \\ a^4x + b^4y + c^4z + d^4u + e^4w &= h^4. \end{aligned}$$

146. Proposed by *G. B. M. ZERR*, A. M., Ph. D., Professor of Chemistry and Physics, The Temple College, Philadelphia, Pa.

Solve by a short original method, if possible:

$$\begin{aligned} x/a + y/b + z/c &= P \dots (1), \\ x/a + b/y + z/c &= Q \dots (2), \\ a/x + y/b + z/c &= R \dots (3). \end{aligned}$$

GEOMETRY.

180. Proposed by *R. TUCKER*, M. A.

ABC is a triangle; A' , B' , C' are the images of A , B , C with respect to BC , CA , AB . The circum-circle ABC cuts $A'BC$ (say) in K (on $A'B$), M (on $A'C$), and AK , AM , AA' cut BC in P , R , Q , respectively. Prove that (1) the orthocenters of the associated triangles lie on circle ABC ; (2) triangle AKM has its sides parallel to and equal twice the sides of the pedal triangle of ABC , and is also equal triangle formed by the above-named orthocenters; (3) $CP \cdot a = b^2$,